

Ceres



Parentage

Ribes dikusche x Barchatnaja

Breeding Center

Instytut Sadownictwa i Kwiaciarnictwa, Skierniewice



Plant growth

short

Plant Habit

spreading

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

?

Antracnose

?

WPBR

?

Gall Mite

?

* *Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

?

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?**acidity**?**anthocyanins**?**ascorbic acids**?

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Ben Gairn



Parentage

Ben Alder x Golubka

Breeding Center

James Hutton Institute, Dundee



Plant growth

short

Plant Habit

upright

Time of flowering

very early

Time of Fruit Ripening

very early

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

?

WPBR

?

Gall Mite

2

* *Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

large (1,11-1,50g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

low **acidity** low **anthocyanins** medium **ascorbic acids** low

* *Ranking scale: very low, low, medium, high, very high*



Best on deep fertile soils, which has never grown currants. Resistant to reversion virus. Attractive to aphids.

Commercial use
Country specifics
other

Ben Dorain



Parentage

Ben Alder x SCRI C2/1/62

Breeding Center

James Hutton Institute, Dundee



Plant growth

tall

Plant Habit

upright

Time of flowering

late

Time of Fruit Ripening

late

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

?

WPBR

?

Gall Mite

7

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?**acidity**low**anthocyanin**sh**high****ascorbic acids**?

** Ranking scale: very low, low, medium, high, very high*



Commercial use
Country specifics
other

Ben Avon



Parentage

Ben Alder x SCRI C2/1/62 (sister seedling of Ben Dorain)

Breeding Center

James Hutton Institute, Dundee



Plant growth

medium

Plant Habit

upright

Time of flowering

late

Time of Fruit Ripening

late

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

?

WPBR

?

Gall Mite

7

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

?

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?acidity?anthocyanins?ascorbic acidshigh

* Ranking scale: very low, low, medium, high, very high



Commercial use
Country specifics
other

Ben Loyal



Parentage

Ben More x Ben Lomond

Breeding Center

James Hutton Institute, Dundee



Plant growth

medium

Plant Habit

upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

?

WPBR

?

Gall Mite

1

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

large (1,11-1,50g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?acidityhighanthocyanins?ascorbic acids?

* *Ranking scale: very low, low, medium, high, very high*



Commercial use
Country specifics
other

Öjebyn



Parentage

Local variety from Norrbotten in North Sweden

Breeding Center

Plant growth

medium

Plant Habit

semi-upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

7

Antracnose

5

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?acidity?anthocyanins?ascorbic acidsmedium

* *Ranking scale: very low, low, medium, high, very high*

N.A.



Commercial use

Country specifics

other

Mortti



Parentage

Öjebyn x Wellington

Breeding Center

?



Plant growth

medium

Plant Habit

upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

7

Antracnose

5

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?acidity?anthocyanins?ascorbic acidsmedium

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

to processing, not good enough for fresh market

Country specifics

other

Ben Connan



Parentage

Ben Sarek x Ben Lomond

Breeding Center

James Hutton Institute, Dundee



Plant growth

medium

Plant Habit

spreading

Time of flowering

early

Time of Fruit Ripening

medium

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

2

Antracnose

2

WPBR

2

Gall Mite

5

* *Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

large (1,11-1,50g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

low **acidity** low **anthocyanins** very low **ascorbic acids** low

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Ben Tron



Parentage

((Vistavotnjaja x (Mendip Cross' x R. dikuscha) x (Goliath x Øjebyn)) x Westra)

Breeding Center

James Hutton Institute, Dundee



Plant growth

tall

Plant Habit

upright

Time of flowering

very early

Time of Fruit Ripening

early

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

2

WPBR

2

Gall Mite

2

* *Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

large (1,11-1,50g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

medium **acidity** low **anthocyanin** high **ascorbic acids** medium

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Ben Lomond



Parentage

(Consort x Magnus) x (Brodtorp x Jaslunda)

Breeding Center

James Hutton Institute, Dundee



Plant growth

medium

Plant Habit

semi-upright

Time of flowering

late

Time of Fruit Ripening

medium

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

9

Antracnose

5

WPBR

5

Gall Mite

7

* *Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

medium **acidity** high **anthocyanin** high **ascorbic acid** high

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Ben Alder



Parentage

Ben More x Ben Lomond

Breeding Center

James Hutton Institute, Dundee



Plant growth

short

Plant Habit

upright

Time of flowering

very late

Time of Fruit Ripening

late

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

9

WPBR

7

Gall Mite

9

* *Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

medium **acidity** medium **anthocyanins** medium **ascorbic acid** high

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other